



Allenton, Wisconsin

Farmers Select Starter R32

MEDICATED

Calves Other Feed

For prevention and control of coccidiosis due to *Eimeria bovis* and *Eimeria zuernii*.

ACTIVE DRUG INGREDIENTS

Monensin	32 gm/ton
----------	-----------

GUARANTEED ANALYSIS

Crude Protein, min %	18.0	Salt (NaCl), min %	0.4
Crude Fat, min %	2.5	Salt (NaCl), max %	0.9
Crude Fiber, max %	8.5	Magnesium (Mg), min %	0.2
ADF, max %	11.5	Selenium (Se), min ppm	0.30
Calcium (Ca), min %	0.6	Vitamin A, min IU/lb	8,760
Calcium (Ca), max %	1.1	Vitamin D, min IU/lb	2,450
Phos (P), min %	0.5	Vitamin E, min IU/lb	112

INGREDIENT STATEMENT

Grain Products, Plant Protein Products, Molasses Products, Roughage Products, Processed Grain By-Products, Calcium Carbonate, Vegetable Oil, Salt, Calcium Phosphate, Monocalcium Phosphate, Potassium Sulfate, Magnesium Sulfate, Natural and Artificial Flavors, Vitamin E Supplement, Vitamin A Supplement, Vitamin D3 Supplement, Sodium Bentonite, Zinc Oxide, Magnesium Oxide, Ferrous Sulfate, Zinc Sulfate, Copper Sulfate, Ethylenediamine Dihydroiodide, Cobalt Carbonate

DIRECTIONS FOR USE

Feed at a rate to provide 0.14 to 1.0 mg/lb body weight up to 200 mg/head/day.

Monensin Level in Feed (gm/ton)	Body Weight (lb)	Minimum lb/head/day	Maximum lb/head/day
32	100	0.88	6.25
	200	1.75	12.5
	300	2.62	12.5
	400	3.5	12.5

Provide clean fresh water at all times.

WARNING. A withdrawal time has not been established for pre-ruminating calves. Do not use in calves to be processed for veal. Do not feed to cattle or other ruminants. Do not feed to sheep due to copper content.

CAUTION. USE AS DIRECTED. Do not allow horses or other equines access to feed containing monensin. Ingestion of monensin by horses has been fatal. Monensin medicated cattle and goat feeds are safe for use in cattle and goats only. Consumption by unapproved species may result in toxic reactions. Do not exceed the levels of monensin recommended in the feeding directions, as reduced average daily gains may result. If feed refusals containing monensin are fed to other groups of cattle, the concentration of monensin in the refusals and amount of refusals fed should be taken into consideration to prevent monensin overdosing.

Manufactured By:
FARMERS GRAIN AND
FEED
528 Railroad Street
Allenton, WI 53002

Product Code: 6112

Net Wgt: 50 lb (22.68 kg)